

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Chromatographic Analysis of Selenium Species

*Aleksandra Sentkowska*

## Abstract

Selenium is an important element in environmental and living organisms that is being essential in very narrow concentration range, while deficiency or toxicity occurs outside this range. However, its toxicity depends not only on its dose but also on its chemical form. In environmental samples, selenium can exist in inorganic forms (as elemental selenium, metal selenides, selenite, or selenate anions) and as organic species with direct C-Se bonds (methylated compounds, selenoaminoacids, and selenoproteins). Thus, the development of reliable techniques to study the speciation of selenium in environmental samples is necessary. The main purpose of this chapter is to provide an update on the recent literature concerning the strategies for selenium speciation in environmental samples. Liquid chromatography coupled with sensitive detector is a commonly used technique for selenium separation. Gas chromatography can also be applied for such purpose; however derivatization step is usually required before analysis. Direct determination of selenium species at the concentration levels present in natural samples is very often difficult or even impossible. For this, several preconcentration/separation procedures for selenium have been proposed, including coprecipitation, extraction into an organic solvent, or application of solid sorbents.

**Keywords:** selenium, speciation, chromatography, gas chromatography, liquid chromatography

## 1. Introduction

Selenium and its several species have been demonstrated to be essential for living organisms [1, 2]. It plays a key role in many important metabolic pathways such as thyroid hormone metabolism and antioxidant defense systems [3]. For these reasons, it should be present in human diet. The range between the deficiency and toxicity of selenium is very narrow. The nutritionally required daily uptake of selenium is 55  $\mu\text{g}$ ; however some studies suggest that it should be 100  $\mu\text{g}$  [4]. It is estimated that the diets of 1 billion people might lack sufficiency for their well-being [5, 6]. The toxic dose of selenium is very much dependent on its chemical form, with different toxicity for organic and inorganic forms [7]. Selenoaminoacids are principal dietary forms of selenium, and selenomethionine (SeMet) is derived from plants, while selenocysteine (SeCys) from animals [8, 9].

Selenium naturally exists in many different inorganic (elemental selenium, selenide, selenite, and selenate ions) and organic forms (methylated compounds, selenoaminoacids, and selenoproteins). Inorganic selenium, present in water and soil, can be easily transformed into volatile compounds by plants and fungi.

	IUPAC name	Abbreviation	Chemical formula
Air	Diethylselenide	DESe	$\text{CH}_3\text{-CH}_2\text{-Se-CH}_2\text{-CH}_3$
Soil and water	Selenite	Se(IV)	$\text{SeO}_3^{2-}$
	Selenate	Se(VI)	$\text{SeO}_4^{2-}$
Plants	Dimethylselenide	DMSe	$\text{CH}_3\text{-Se-CH}_3$
	Dimethyldiselenide	DMDSe	$\text{CH}_3\text{-Se-Se-CH}_3$
	Dimethylseleniumsulfide	DMSSe	$\text{CH}_3\text{-Se-S-CH}_3$
	Selenocysteine	SeCys	$\text{HSe-CH}_2\text{CH(NH}_2\text{)-COOH}$
	Selenocystine	SeCys <sub>2</sub>	$\text{HCOOH-CH(NH}_2\text{)CH}_2\text{-Se-Se-CH}_2\text{CH(NH}_2\text{)-COOH}$
	Se-Methylselenocysteine	SeMC	$\text{CH}_3\text{Se-CH}_2\text{CH(NH}_2\text{)-COOH}$
	Selenomethionine	SeMet	$\text{CH}_3\text{Se-CH}_2\text{CH}_2\text{CH(NH}_2\text{)-COOH}$
	Se-Methylselenomethionine	MeSeMet	$(\text{CH}_2)_2\text{Se}^+\text{-CH}_2\text{-CH}_2\text{-CH(NH}_2\text{)-COOH}$
	$\gamma$ -Glutamyl-Se-methylselenocysteine	$\gamma$ -Glu-SeMC	$\text{H}_2\text{NCH}_2\text{CH}_2\text{-CO-NGCH(COOH)CH}_2\text{-Se-CH}_3$
	Selenocystathionine	SeCysth	$\text{HCOOH-CH(NH}_2\text{)-CH}_2\text{-CH}_2\text{-Se-CH}_2\text{-CH(NH}_2\text{)COOH}$
	Selenocystamine	SeCystm	$\text{H}_2\text{N-CH}_2\text{-CH}_2\text{-Se-Se-CH}_2\text{-CH}_2\text{-NH}_2$
	Se-Adenosylselenohomcysteine	AdoSeHcy	$\text{NH}_2\text{-CH(COOH)-CH}_2\text{-CH}_2\text{-Se-CH}_2\text{-C}_4\text{H}_5\text{O}_3\text{C}_5\text{N}_4\text{-NH}_2$

**Table 1.**  
*Principal selenium species present in environmental samples.*

Organic species of selenium form covalent C-Se bonds. SeCys is included into selenoproteins and participates in redox reactions. The metabolic pathway of selenium in the human body is complicated [10]. In general it can be divided into three groups including reduction of inorganic species by glutathione (GHS) to selenide, cleavage reaction of organic species by  $\beta$ -lyase; utilization according to the UGA codon leading the synthesis of selenoproteins; and finally excretion after being metabolized to methylated species.

Apart from the importance of the selenium in living organisms, this element is also spread throughout the environment. Sulfur-containing minerals are natural sources of selenium, but it is also produced by combustion of fossil fuels. It should be noticed that selenium is used in electronic industries and agriculture as a component of fertilizers.

Several analytical procedures for determination of selenium at low concentration levels in environmental samples have been proposed and recently reviewed [11, 12]. For selenium speciation analysis, the coupling of chromatographic techniques such as gas chromatography (GC) or high-performance liquid chromatography (HPLC) with a highly sensitive and selective detector is very useful [13]. Even though GC exhibits high efficiency and simplicity, HPLC has the ability of dealing with non-volatile compounds, extending the range of application and avoiding a derivatization step. This chapter will focus on the recent progress in the application of HPLC in different modes for selenium speciation analysis in water, soil, and plants. Sample pretreatment procedures will be also considered. The principal selenium species present in environmental samples are summarized in **Table 1**.

## 2. Sample preparation

Sample preparation step is crucial in every analysis where analytes are present at very low concentration levels. In the speciation analysis, there is also another difficulty that should be overcome. An important requirement for reliable speciation is to retain the concentration and structure of the original chemical forms in the sample. In general, aquatic samples such as rain, ground- or surface water, tap and drinking water, seawater, and soil solutions do not require any pretreatment procedures other than filtration through 0.45  $\mu$ m filter.

Extraction of selenium species from the solid samples with the highest recovery is quite challenging. According to Peachey et al. [14], selection of the extraction method, which provides high extraction efficiency while preventing the integrity of selenium species, is essential for the accurate measurement of its species. The most used method can be divided into three main groups:

- Extraction using aqueous solution (water, water-methanol, and buffers).
- Acid or alkali hydrolysis (hydrochloric acid and methanesulfonic acid).
- Enzymatic hydrolysis (proteinase, protease, and mixture of enzymes).

Since selenoaminoacids are water-soluble, extraction with hot water is extensively used [14–17]. However, the efficiency of water extraction from yeast was only 10% [17]. To release bounded selenoaminoacids, enzymatic or acidic hydrolysis was necessary [18–20]. The addition of methanesulfonic acid was used for selenomethionine extraction from yeast when heated under reflux [21]. Casiot et al. [22] reported that extraction of selenium species from yeast with water and ethanol led only to 10–20% recoveries of selenium and not allowed to extract

selenomethionine. The addition of pectinolytic enzymes released additional 20% of selenomethionine, while the addition of dodecyl sulfate solution allowed solubilization of a selenoprotein that accounted for 30% of total selenium. On the other hand, using tetramethylammonium hydroxide solubilized the sample completely, but the extracted selenium species were entirely degraded to selenomethionine and inorganic selenium [22]. It should be noticed that this type of extraction strictly depends on the choice of an enzyme, pH of the extraction solution, as well as temperature and time of the extraction. The most commonly used enzymes for this purpose are proteinase K or proteolytic enzymes (protease XIV), which were used for water-insoluble selenium fraction in many complicated matrices [23]. To reduce the extraction time, also ultrasonic hydrolysis can be used as the breakdown of Se-containing proteins (peptide bonds) into selenoaminoacids occurs [24]. The *in vitro* digestion with gastric juice was also used for selenium extraction from fish samples [25]. This procedure allowed SeMet determination, but the whole process takes few hours.

The sequential procedure developed by Chassaigne et al. [26] consisted of three steps: first, Tris-HCl buffer was used for extraction of water-soluble fractions, then also Tris-HCl with the addition of SDS for solubilization of protein fraction, and, finally, concentrated HNO<sub>3</sub> was used for dissolving of the remaining solid residue. A similar three-step procedure was used for extraction of selenium from mushrooms with 89% extraction efficiency [27]. It should be noticed that sequential treatment was also applied for dietary supplements [28, 29].

Cuderman et al. [30] examined different extraction media to identify selenium species in buckwheat. The optimal extraction efficiencies were obtained by hydrolysis with HCl, followed by breaking the cells with liquid nitrogen and then enzymatic hydrolysis with protease XIV. Ammonium acetate [31], sodium hydroxide [32], and enzymatic hydrolysis with pronase E [33] have been proposed for extraction of selenium species from green onion leaves.

### **3. Total selenium determination**

Determination of the total selenium content is still the first step of its analysis. This procedure requires that organic forms must be transferred into inorganic selenium that is usually achieved with digestion using strong mineral acids or UV irradiation after addition of hydrogen peroxide [12, 34]. For this purpose, fluorometry, electrochemical detectors, atomic absorption spectrometry (AAS), atomic fluorescence spectrometry (AFS), mass spectrometry (MS), and neutron activation analysis have been used (Table 2).

Hydride generation coupled to AAS or AFS detectors is specific for Se(IV) determination, where these species are selectively reduced to volatile SeH<sub>2</sub>, usually by sodium tetrahydroborate in hydrochloric medium. This technique can be applied for the determination of total inorganic selenium (e.g., sum of selenite and selenate) after quantitative reduction. The content of Se(VI) is then obtained by the difference between two determinations. This technique can be fully automatic by connection with flow injection analysis system. The advantage of such system is minimum sample and reagent consumption as well as short time of single run.

Stripeikis et al. [44] determined selenite and selenate in drinking water using fully automatic online separation/pre-concentration system coupled to electrochemical atomic spectrometry. Preconcentration of both selenium forms was carried out onto a microcolumn packed with an anionic resin (Dowex 1X8) that was placed in the robotic arm of the autosampling device. Selenite and selenate were then sequentially eluted with HCl at concentration of 0.1 and 4 mol L<sup>-1</sup>,



Technique	Matrix	Detection limit	Ref.
UV-Vis	Water, soil	0.012 mg L <sup>-1</sup>	[34]
Fluorometry	Water	0.35 ng	[35]
X-ray fluorescence	Water	0.032 ng mL <sup>-1</sup>	[36]
HGAAS	Water	2 ng L <sup>-1</sup>	[37]
ETAAS	Seafood	0.16 µg g <sup>-1</sup>	[38]
ICP-MS	Soil	3–29 ng L <sup>-1</sup>	[39]
Adsorptive stripping voltammetry	Water	0.06 ng mL <sup>-1</sup>	[40]
Differential-pulse cathodic stripping voltammetry	Food supplements	0.2 ng mL <sup>-1</sup>	[41]
Differential-pulse anodic stripping voltammetry		0.06 ng mL <sup>-1</sup>	[42]
Instrumental neutron activation analysis	Libyan food	26–90 µg L <sup>-1</sup>	[43]

**Table 2.**  
*Analytical methods for the determination of total selenium.*

respectively. The interference of large amounts of chloride ions during selenium atomization was prevented by using iridium as a permanent chemical modifier.

Kocot et al. [36] proposed a dispersive micro-solid-phase extraction with graphene as a solid adsorbent and ammonium pyrrolidine dithiocarbamate as a chelating agent for Se(IV) in analysis of inorganic selenium by the energy-dispersive X-ray fluorescence spectrometry; the concentration of Se(VI) was calculated as the difference between the concentration of selenite after and before reduction of selenate.

Due to the high selectivity and sensitivity, wide linear range, as well as multielement and multi-isotope detection, inductively coupled plasma mass spectrometry (ICP-MS) is a great tool for selenium analysis. However, some difficulties can be found when conventional ICP-MS is used for this purpose. To avoid the spectra interferences with <sup>80</sup>Se isotope (the largest natural abundance of 49.6%), <sup>82</sup>Se and <sup>77</sup>Se are often monitored. The use of collision/reaction cell, operating with hydrogen gas, lowers argon dimer interferences. This technique was applied by Reyes et al. to determine the selenium in biological reference materials and serum samples [45] and yeast [46].

#### 4. Separation techniques in selenium speciation analysis

The occurrence of selenium at very low concentration levels as well as the dependence of its toxicity on the form in which it occurs resulted in the need for reliable analytical procedures for identification and quantification of its species. The coupling of separation of selenium species with sensitive detection has become a powerful technique for Se speciation. Liquid chromatography operating in different modes is the most used analytical technique for this purpose; however electrophoresis and gas chromatography also were used in selenium analysis. That is why these two methods will be also described in this chapter, which will mainly focus on liquid chromatography.

##### 4.1 Gas chromatography

Selenium species can be divided into volatile and nonvolatile compounds. The first group of compounds can be directly analyzed using gas chromatography,

for example, dimethylselenide was determined in human breath after ingestion of Se-enriched selenite [47]. For other nonvolatile compounds, derivatization is required.

In the literature many derivatization procedures can be found. Pelaez et al. [48] tested two methods of derivatization of selenomethionine and selenomethionine in selenium supplements. The first method consisted of esterification of the carboxylic acid group using propan-2-ol and then acylation of amino group. The second described procedure used ethanol-ethyl chloroformate for one step derivatization. Then detection using ICP-MS was performed.

In general, many types of detectors were coupled with GC for speciation analysis of selenium. Yang et al. [49] successfully used ICP detection for the determination of selenium in yeast. Organoselenium species in plants were determined using GC-MIP-AES detection [50]. Because of the requirement of the derivatization, GC is not so widely used in selenium speciation analysis in comparison to liquid chromatography.

## **4.2 Capillary electrophoresis**

Due to the high resolving power, capillary electrophoresis (CE) has a potential to be used in speciation analysis as an alternative or complementary technique to HPLC. CE coupled with ICP-MS was used in the analysis of selenium in yeast extract after SEC separation [51]. The obtained limit of detection for organic and inorganic species was in range 7–18  $\mu\text{g L}^{-1}$ . The main difficulty in such separation can be big sizes of selenopolypeptides resulting in their slow migration. In such case, the predigestion of selenoprotein fraction is recommended.

It should be highlighted that gel electrophoresis offers better resolution than liquid chromatography in the analysis of high molecular weight selenoproteins, which was used in such analysis by Chassaing et al. [26], and in Se-enriched yeast analysis as well by Chery et al. in blood sample analysis [52].

## **4.3 High-performance liquid chromatography**

### *4.3.1 Size-exclusion chromatography*

Separation in size-exclusion chromatography (SEC) strictly depends on the size of separated analytes. That is why this technique is widely used as a preliminary step to sample purification or to separate selenoproteins from the matrix. Ayouni [53] observed high molecular weight selenium compounds in the extract of dietary supplements after separation by SEC. Conjunction of SEC with ICP-MS was used for analysis of the products of enzymatic digestion of selenoproteins fraction [54, 55]. Size-exclusion chromatography was also used in selenium analysis in human plasma [56] and extracts from rats' internal organs [57].

### *4.3.2 Ion-exchange chromatography*

Anion-exchange chromatography has been mainly employed in selenium speciation analysis [58–63]. Mobile phases used in this mode usually contain small content of organic modifier (e.g., 2–5% methanol) and buffered salt solution (e.g., acetate, phosphate, and citrate). During the separation process, the equilibria between the charged solute ion and the oppositely charged surface of the stationary phase are established. The separation is achieved based on the differences of the strength of such interaction between analytes. Anion-exchange

chromatography was used in the selenium speciation analysis in garlic, sunflower, and radish sprouts [64]. In addition to well-known compounds like SeMet or MeSeCys, several unidentified signals were obtained. The application of high-resolution mass spectrometry enabled identification of additional seleno-compounds as inorganic metabolites, such as deamino-hydroxy-seleno-homolanthionine, *N*-acetylcysteine-selenomethionine, methylseleno-pentose-hexose, methylselenoglutathione, 2,3-dihydroxy-propionyl selenocysteine-cysteine, methyltio-selenoglutathione, 2,3-dihydroxy propionyl-seleno-lanthionine, and two Se-containing compounds with proposed molecular formula  $C_{10}H_{18}N_2O_6Se$  and  $C_{10}H_{13}N_5O_3Se$ .

Cation-exchange chromatography was used to analyze selenium-enriched yeast in a human adsorption study [65]. As a mobile phase, pyridinium formate buffer with 3% of methanol was used. This method was suitable for separation of organic selenium species, however not suitable to separate selenite and selenate.

#### *4.3.3 Reversed-phase chromatography*

Both, simple reversed-phase [66–68] and ion-pair (IP) reversed-phase chromatography [15, 28, 69], are widely used for analysis of ionic and neutral selenium species. The mobile phases are aqueous with small amount of polar organic solvent (usually methanol or acetonitrile). Because of their hydrophilicity, selenoaminoacids are not retained onto typical reversed-phase columns. The use of ion-pairing reagents as mobile phase additives allows their separation. The ion-pairing reagent is usually an alkyl sulfonate, an alkyl sulfate, or an alkylammonium salt. Its non-polar chain interacts with hydrophobic stationary phase (e.g., C8 or C18), while ionizable group is neutralized by oppositely charged analyte. Hexanesulfonic acid has been used as anion-pairing reagent in the speciation analysis of selenium in Brazil nuts, using C8 column for separation [70]. Obtaining separation was satisfied for organic compounds but poor for selenite and selenate. For separation of organic and inorganic forms of selenium, tetrabutylammonium acetate was proposed [71]. Also mixed ion-pairing reagents (butanesulfonic and tetramethylammonium hydroxide) were also used to simultaneously separate inorganic and organic species with satisfactory separation efficiencies [72].

New mobile phase additives are still developed, for example, room temperature ionic liquids [73]. Their mechanism of action is based on bilayer formulation onto stationary phase. It gives the possibility of additional interactions between the analyte and the bed, which significantly affects the retention and shape of the recorded signals. The effects of several imidazolium chlorides on the separation of selenium species mixture was described in details [74]. In all cases, SeMeCys was the first species eluted indicating its weak retention in the column, while the retention time of Se(VI) was increasing with the increase of alkyl chain.

#### *4.3.4 Hydrophilic interaction liquid chromatography*


Hydrophilic interaction liquid chromatography (HILIC) is a complementary technique to reversed-phase mode. The separation mechanism is mainly based on the partition of the analyte between the thin water layer adsorbed onto the stationary surface and the eluent, which contain high content of organic solvent usually acetonitrile. It is known that also other interactions such as hydrogen bonding dipole-dipole interactions and electrostatic forces may play an important role in the retention mechanism in HILIC [75]. The governed retention mechanism strictly depends on the type of used stationary phase and the buffer conditions




(content of organic solvent, concentration of salt, and pH). TSKgel Amide-80 stationary phase with covalently bound carbamoyl groups is frequently used in the analysis of selenium in HILIC mode [76, 77]. According to the characterization of amide stationary phase, the achieved separation is not pH dependent [78]. Also zwitterionic and silica stationary phases have been also used in HILIC separation of selenium [79, 80]. It should be highlighted that in selenium separation in HILIC mode methanol is recommended instead of acetonitrile as a main component of mobile phase [79] which is shown in **Figure 1**.

The use of methanol enhances peak intensity, improve the separation of SeMet and MeSeCys, and shorten time of the single run. The best separation conditions

**A**



**B**




**Figure 1.**

The chromatogram of selenium compounds obtained on silica column using (A) MeOH and (B) ACN in the mobile phase [79]. Reprinted with permission from Elsevier.

have been obtained for silica column and mobile phase consisted of 85% of methanol and 8 mM of ammonium acetate. Using the zwitterionic column (ZIC-HILIC) instead of silica stationary phase resulted in recording of very asymmetric peaks.

The potential of two orthogonal chromatographic modes—RP and HILIC—was examined in the analysis of onion leaf extracts [79]. Higher separation efficiency (mainly for inorganic selenium species) and shorter retention times were obtained when HILIC mode was used (**Figure 2**).



**Figure 2.** The chromatographic separation of selenium species present in onion leaf extracts in (A) HILIC mode and (B) RP mode. (A) Atlantis HILIC (silica) column, mobile phase: 85% MeOH and 8 mM  $\text{CH}_3\text{COONH}_4$ , pH 7. (B) Luna C8 column, mobile phase: 99.5%  $\text{HCOOH}$  and 0.5% ACN [79]. Reprinted with permission from Elsevier.

5. Conclusions

The determination of selenium is of great importance from the point of view of understanding its metabolism and its potential benefits for human health. Due to the variety of selenium species and low concentrations in which it occurs in environmental samples, liquid chromatography coupled to ICP-MS is the most powerful method for speciation analysis of selenium. However, other techniques, for example, HILIC-MS/MS, have also been successfully applied in the speciation analysis of selenium. It should be highlighted that for such analysis all the analytical steps like selection of separation and detection method and the optimization of separation parameters (eluent type and composition, mobile phase additives) should be optimized to avoid the coelution of selenium species.

The recent application of liquid chromatography for selenium separation in plants (vegetables samples) is shown in **Table 3**. For sure new chromatographic strategies will be described in the literature in the nearest future.

Sample (determined selenium species)	Mobile phase	Stationary phase	Detection	Ref.
Reversed-phase chromatography				
Green leafy vegetables (SeCys <sub>2</sub> , SeMet)	50% MeOH + 1.5% HCOOH	C18 (250 × 4.6 mm, 5 μm)	ICP-MS	[69]
Ion-pair reversed-phase chromatography				
Garlic (MeSeCys, SeMet, Se(IV), Se(VI))	Gradient phosphate buffer pH 6.0, 5% MeOH, 0.1% [C6mim]Cl	Agilent Zorbax SB-C8 (150 × 4.6 mm)	HG-AFS	[58]
Anion-exchange chromatography				
Onion (MeSeCys, SeMet, Se(IV), g-glu-MeSeCys, Se(VI))	Gradient acetate buffer, pH 4.7	Hamilton PRP-X100 (250 × 4.1 mm, 10 mm)	ICP-MS	[69]
Cation-exchange chromatography				
Carrot (inorganic Se, SeMet, g-glu-MeSeCys)	Gradient: ammonium formate, pH 3.0, 5% MeOH	Chrompack IonoSpher 5C (150 × 2.0 mm, 5 mm)	ICP-MS	[75]
Hydrophilic liquid chromatography				
Onion leaves (Se(IV), Se(VI), MeSeCys, SeMet)	Gradient: MeOH and ammonium acetate, pH 7	Atlantis HILIC (silica) (100 × 2.1 mm, 3.0 mm)	MS/MS	[81]

**Table 3.**  
*Examples of the recent HPLC application for selenium speciation in plant materials.*

Conflict of interest

The author declares no conflict of interest.

Appendices and nomenclature

- AAS
- AFS
- atomic absorption spectrometry
- atomic fluorescence spectrometry

ACN	acetonitrile
C8	octylsilane column
CE	capillary electrophoresis
CH <sub>3</sub> COONH <sub>4</sub>	ammonium acetate
ETAAS	electrothermal atomic absorption
GC	gas chromatography
HCOOH	formic acid
HCl	hydrochloric acid
HGAAS	hydride generation atomic absorption spectrometric
HILIC	hydrophilic interaction liquid chromatography
HNO <sub>3</sub>	nitric acid
HPLC	high-performance liquid chromatography
ICP-MS	inductively coupled plasma-mass spectrometry
ILs	ionic liquids
IP	ion pair
MeOH	methanol
MeSeMet	Se-methylselenomethionine
RP	reverse phase
Se(IV)	selenite
Se(VI)	selenate
SeCys	selenocysteine
SEC	size-exclusion chromatography
SeMet	selenomethionine

### Author details

Aleksandra Sentkowska  
Heavy Ion Laboratory, University of Warsaw, Warsaw, Poland

\*Address all correspondence to: [sentkowska@slcj.uw.edu.pl](mailto:sentkowska@slcj.uw.edu.pl)

### IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Tan LC, Nancharaiach YV, van Hullebusch ED, Lens PNL. Selenium: Environmental significance, pollution and biological treatment technologies. *Biotechnology Advances*. 2016;**34**:886-907. DOI: 10.1016/j.biotechadv.2016.05.005
- [2] Weekeley CM, Harris HH. Which form is that? The importance of selenium speciation and metabolism in the prevention and treatment of disease. *Chemical Society Reviews*. 2013;**42**:8870-8894. DOI: 10.1039/c3cs60272a
- [3] Zimmerm MT, Bayse CA, Ramoutar RR, Brumaghim JL. Sulfur and selenium antioxidants: Challenging radicals scavenging mechanism and developing structure-activity relationship based on metal binding. *Journal of Inorganic Biochemistry*. 2015;**145**:30-40. DOI: 10.1016/j.jinorgbio.2014.12.020
- [4] Clark LC, Combs GF, Turnbull BW, Slate EH, Chalker DK, Chow J, et al. Effects of selenium supplementation for cancer prevention in patients with carcinoma of the skin. A randomized controlled trial. Nutritional Prevention of Cancer Study Group. *Journal of the American Medical Association*. 1996;**276**:1957-1963
- [5] Joy EJM, Ander EL, Young SD, Black CR, Watts MJ, Chilimba ADC, et al. Dietary mineral supplies in Africa. *Physiologia Plantarum*. 2014;**31**:208-229. DOI: 10.1111/ppl.12144
- [6] Stoffaneller R, Morse NL. A review of dietary selenium intake and selenium status in Europe and the Middle East. *Nutrients*. 2015;**7**:1494-1537. DOI: 10.3390/nu7031494
- [7] Forchhammer K, Bock A. Biologie und biochemie des elements selen. *Naturwissenschaften*. 1991;**78**:497-504. DOI: 10.1007/BF01131397
- [8] Olsen OE, Novacek EJ, Whitehead EI, Palmer IS. Investigation of selenium in wheat. *Phytochemistry*. 1970;**9**:1181-1188. DOI: 10.1016/S0031-9422(00)85306-6
- [9] Hawkens WC, Wilhelmes EC, Tapper AL. Abundance and tissue distribution of selenocysteine-containing proteins in the rat. *Journal of Inorganic Biochemistry*. 1985;**23**:77-92. DOI: 10.1016/0162-0134(85)83011-7
- [10] Wastney ME, Combs GF, Canfield WK, Taylor PR, Patterson KY, Hill DA, et al. A human model of selenium that integrates metabolism from selenite and selenomethionine. *Japanese Journal of Nutrition*. 2011;**141**:708-717. DOI: 10.3945/jn.110.129049
- [11] Kumkrong P, LeBlanc K, Mercier PHJ, Mester Z. Selenium analysis in waters. Part 1: Regulation and standard methods. *Science of the Total Environment*. 2018;**640-641**:1611-1634. DOI: 10.1016/j.scitotenv.2018.05.392
- [12] LeBlanc K, Kumkrong P, Mercier PHJ, Mester Z. Selenium analysis in waters. Part 2: Speciation methods. *Science of the Total Environment*. 2018;**640-641**:1635-1651. DOI: 10.1016/j.scitotenv.2018.05.394
- [13] Marcinkowska M, Barańkiewicz D. Multielemental speciation analysis by advanced hyphenated technique—HPLC/ICP-MS: A review. *Talanta*. 2016;**161**:1777-1204. DOI: 10.1016/j.talanta.2016.08.034
- [14] Peachey P, McCarthy N, Goenaga-Infante H. Acceleration of enzymatic hydrolysis of protein-bound selenium by focused microwave energy. *Journal of Analytical Atomic Spectrometry*. 2008;**23**:487-492. DOI: 10.1039/B717854A
- [15] Kubacha KM, Hanley T, Mantha M, Wiklson RA, Falconer TM, Kassa Z,



et al. Evaluation of selenium in dietary supplements using elemental speciation. *Food Chemistry*. 2017;**218**:313-320. DOI: 10.1016/j.foodchem.2016.08.086

[16] Tsai Y, Lin C, Hsu I, Sun Y. Sequential photocatalyst-assisted digestion and vapor generation device coupled with anion exchange chromatography and inductively coupled plasma mass spectrometry for speciation analysis of selenium species in biological samples. *Analytica Chimica Acta*. 2014;**806**:165-171. DOI: 10.1016/j.aca.2013.11.008

[17] Kotrebai M, Bird SM, Tyson JF, Block E, Uden PC. Characterization of selenium species in biological extracts by enhanced ion-pair liquid chromatography with inductively coupled plasma-mass spectrometry and by referenced electrospray ionization-mass spectrometry. *Spectrochimica Acta B*. 1999;**54**:1573-1591. DOI: 10.1016/S0584-8547(99)00086-5

[18] Kotrebai M, Birringer M, Tyson JF, Block E, Uden PC. Selenium speciation in enriched and natural samples by HPLC-ICP-MS and HPLC-ESI-MS with perfluorinated carboxylic acid ion-pairing agents. *Analyst*. 2000;**125**:71-78. DOI: 10.1039/A906320J

[19] Kotrebai M, Tyson JF, Block E, Uden PC. High-performance liquid chromatography of selenium compounds utilizing perfluorinated carboxylic acid ion-pairing agents and inductively coupled plasma and electrospray ionization mass spectrometric detection. *Journal of Chromatography. A*. 2000;**866**:51-63. DOI: 10.1016/S0021-9673(99)01060-2

[20] McSheehy S, Yang W, Pannier F, Szpunar J, Łobiński R, Auder J, et al. Speciation analysis of selenium in garlic by two-dimensional high-performance liquid chromatography with parallel inductively coupled plasma mass

spectrometric and electrospray tandem mass spectrometric detection. *Analytica Chimica Acta*. 2000;**421**:147-153. DOI: 10.1016/S0003-2670(00)01039-4

[21] Wróbel K, Wróbel K, Caruso JA. Selenium speciation in low molecular weight fraction of Se-enriched yeast by HPLC-ICP-MS: Detection of selenoadenosylmethionine. *Journal of Analytical Atomic Spectrometry*. 2002;**17**:1048-1054. DOI: 10.1039/B200920J

[22] Casiot C, Szpunar J, Łobiński R, Potin-Gautier M. Sample preparation and HPLC separation approaches to speciation analysis of selenium in yeast by ICP-MS. *Journal of Analytical Atomic Spectrometry*. 1999;**14**:645-650

[23] Huerta VD, Sanchez MLF, Sanz-Medel A. Quantitative selenium speciation in cod muscle by isotope dilution ICP-MS with reaction cell: Comparison of different reported extraction procedures. *Journal of Analytical Atomic Spectrometry*. 2004;**19**:644-648. DOI: 10.1039/b313826g

[24] Capelo JL, Fernandez C, Pedras B, Santos P, Gonzales P, Vaz C. Trends in selenium determination/speciation by hyphenated techniques based on AAS or AFS. *Talanta*. 2006;**68**:1442-1447. DOI: 10.1016/j.talanta.2005.08.027

[25] Cabanero AI, Madrid Y, Camara C. Selenium and mercury bioaccessibility in fish samples: An in vitro digestion method. *Analytica Chimica Acta*. 2004;**526**:51-61. DOI: 10.1016/j.aca.2004.09.039

[26] Chassaigne H, Chery CC, Bordin G, Ropdriguez AR. Development of new analytical methods for selenium speciation in selenium-enriched yeast material. *Journal of Chromatography. A*. 2001;**976**:409-422. DOI: 10.1016/S0021-9673(02)00945-7

- [27] Dernovics M, Stefanka Z, Fodor P. Improving selenium extraction by sequential enzymatic processes for Se-speciation of selenium-enriched *Agaricus bisporus*. Analytical and Bioanalytical Chemistry. 2002;**372**:473-480. DOI: 10.1007/s00216-001-1215-5
- [28] Hsieh Y, Jiang S. Determination of selenium compounds in food supplements using reversed-phase liquid chromatography-inductively coupled plasma mass spectrometry. Microchemical Journal. 2013;**110**:1-7. DOI: 10.1016/j.microc.2013.01.009
- [29] Wang W, Chen Z. Extraction of selenium species in pharmaceutical tablets using enzymatic and chemical methods. Microchimica Acta. 2009;**165**:167-172. DOI: 10.1007/s00604-008-0115-1
- [30] Cuderman P, Ozbolt L, Krefl I, Stibilij V. Extraction of Se species in buckwheat sprouts grown from seeds soaked in various Se solutions. Food Chemistry. 2010;**123**:941-948. DOI: 10.1016/j.foodchem.2010.04.063
- [31] Tie M, Li B, Sun T, Guan W, Liang Y, Li H. HPLC-ICP-MS speciation of selenium in Se-cultivated *Flammulina velutipes*. Arabian Journal of Chemistry. 2017 (in press). DOI: 10.1061/j.arabje.2017.05.012
- [32] Kapolna E, Fodor P. Speciation analysis of selenium enriched green onions (*Allium fistulosum*) by HPLC-ICP-MS. Microchemical Journal. 2006;**84**:52-62. DOI: 10.1016/j.microc.2006.04.014
- [33] Kapolna E, Shah M, Caruso JA, Fodor P. Selenium speciation studies in Se-enriched chives (*Allium schoenoprasum*) by HPLC-ICP-MS. Food Chemistry. 2007;**101**:1398-1406. DOI: 10.1016/j.foodchem.2006.03.048
- [34] Ramachandran Kumar GS. Modified spectrophotometric method for the determination of selenium in environmental and mineral mixtures using 2,3-diaminonaphthalene. Talanta. 1996;**43**:1711-1714. DOI: 10.1016/0039-9140(96)01947-9
- [35] Wang D, Alfthan G, Aro A. Determination of total selenium and dissolved selenium species in natural waters by fluorometry. Environmental Science and Technology. 1994;**28**:383-387. DOI: 10.1021/es00052a007
- [36] Kocot K, Leardi R, Walczak B, Sitko R. Determination and speciation of trace and ultratrace selenium ions by energy-dispersive X-ray fluorescence spectrometry using graphene as solid adsorbent in dispersive micro-solid phase extraction. Talanta. 2015;**134**:360-365. DOI: 10.1016/j.talanta.2014.11.036
- [37] Örnemark U, Pettersson J, Olin A. Determination of total selenium in water by atomic-absorption spectrometry after hydride generation and preconcentration in a cold trap system. Talanta. 1992;**39**:1089-1096. DOI: 10.1016/0039-9140(92)80206-S
- [38] Méndez H, Alava F, Lavilla I, Bendicho C. Ultrasonic extraction combined with fast furnace analysis as an improved methodology for total selenium determination in seafood by electrothermal-atomic absorption spectrometry. Analytica Chimica Acta. 2002;**452**:217-222. DOI: 10.1016/S0003-2670(01)01475-1
- [39] Tolu J, Hecho IL, Bueno M, Thiry Y, Potin-Gautier M. Selenium speciation analysis at trace level in solis. Analytica Chimica Acta. 2011;**684**:126-133. DOI: 10.1016/j.aca.2010.10.044
- [40] Ashournia M, Aliakbar A. Determination of Se(IV) in natural waters by adsorptive stripping voltammetry of 5-nitropiazselenol. Journal of Hazardous Materials.

2010;**174**:788-794. DOI: 10.1016/j.jhazmat.2009.09.121

[41] Holak W, Specchio JJ. Determination of selenium in food supplements by differential-pulse cathodic stripping voltammetry in the presence of added copper. *Analyst*. 1994;**119**:2179-2182. DOI: 10.1039/AN9941902179

[42] Ramadan AA, Mandil H, Shikh-Debes A. Differential pulse anodic stripping voltammetric determination of selenium (IV) at a gold electrode modified with 3, 3'-diaminobenzidine. 4HCl-Nafion. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2014;**6**:148-153. DOI: 10.24297/jac.v15i2.7818

[43] Elghawi U, Al-Sadeq AA, Bejey MM, Alamin MB. Determination of selenium in Libyan food items using pseudocyclic instrumental neutron activation analysis. *Biological Trace Element Research*. 2005;**107**:61-72. DOI: 10.1385/BTER:107:1:061

[44] Stripeikis J, Pedro J, Bonivardi A, Tudino M. Determination of selenite and selenite in drinking water: A fully automatic on-line separation/pre-concentration system coupled to electrochemical atomic spectrometry with permanent chemical modifiers. *Analytica Chimica Acta*. 2004;**502**:99-105. DOI: 10.1016/j.aca.2003.09.022

[45] Reyes LH, Gayon JMM, Alonso JIG, Sanz-Medel A. Determination of selenium in biological materials by isotope dilution analysis with an octopole reaction system ICP-MS. *Journal of Analytical Atomic Spectrometry*. 2003;**18**:11-16. DOI: 10.1039/B209213A

[46] Reyes LH, Marchante-Gayon JM, Alonso JIG, Sanz-Medel A. Application of isotope dilution analysis for evaluation of extraction conditions in the determination of total selenium

and selenomethionine in yeast-based nutritional supplements. *Journal of Agricultural and Food Chemistry*. 2006;**54**:1557-1563. DOI: 10.1021/jf0523768

[47] Kremer D, Ilgen G, Feldman J. GC-ICP-MS determination of dimethylselenide in human breath after ingestion of 77 Se-enriched selenite: Monitoring of in-vivo methylation of selenium. *Analytical and Bioanalytical Chemistry*. 2005;**383**:509-515. DOI: 10.1007/s00216-005-0001-1

[48] Pelaez MV, Bayon MM, Alonso JIG, Sanz-Medel A. A comparison of different derivatisation approaches for the determination of selenomethionine by GC-ICP-MS. *Journal of Analytical Atomic Spectrometry*. 2000;**15**:1217-1222. DOI: 10.1039/B000505N

[49] Yang L, Mester Z, Sturgeon RE. Determination of methionine and selenomethionine in yeast by specie-specific isotope dilution GC/MS. *Analytical Chemistry*. 2004;**75**:5149-5156. DOI: 10.1021/ac049475p

[50] Kahakachchi C, Boakye HT, Uden PC, Tyson JF. Chromatographic speciation of anionic and neutral selenium compounds in Se-accumulating *Brassica juncea* (Indian mustard) and selenized yeast. *Journal of Chromatography. A*. 2004;**1054**:303-312. DOI: 10.1016/j.chroma.2004.07.083

[51] Mounicou S, McSheehy S, Szpunar J, Potin-Gautier M, Łobiński R. Analysis of selenium yeast for selenium speciation by size-exclusion chromatography and capillary zone electrophoresis with inductively coupled plasma mass spectrometric detection (SEC-CZE-ICP-MS). *Journal of Analytical Atomic Spectrometry*. 2002;**17**:15-20. DOI: 10.1039/B107943N



- [52] Chery CC, Gunther D, Cornelis R, Vanhaecke F, Moens L. Detection of metals in proteins by means of polyacrylamide gel electrophoresis and laser ablation-inductively coupled plasma-mass spectrometry: Application to selenium. Electrophoresis. 2003;**24**:3305-3313. DOI: 10.1002/elps.200305590
- [53] Ayouni L. Speciation of selenium in a commercial dietary supplement by liquid chromatography coupled with inductively coupled plasma-mass spectrometry (ICP-MS). Journal of Toxicology and Environmental Health, Part A. 2007;**70**:735-741. DOI: 10.1080/15287390701236314
- [54] Moreno P, Quijano MA, Gutierrez AM, Perez-Conde MC, Camara C. Study of selenium species distribution in biological tissues by size exclusion and ion exchange chromatography inductively coupled plasma-mass spectrometry. Analytica Chimica Acta. 2004;**524**:315-327. DOI: 10.1016/j.aca.2004.02.029
- [55] Siwek M, Galunsky B, Niemeyer B. Isolation of selenium organic species from arctic krill after enzymatic hydrolysis. Analytical and Bioanalytical Chemistry. 2005;**281**:737-741. DOI: 10.1007/s00216-004-2936-z
- [56] Koyama H, Omura K, Ejima A, Kasanuma Y, Watanabe C, Satoh H. Separation of selenium-containing proteins in human and mouse plasma using tandem high-performance liquid chromatography columns coupled with inductively coupled plasma mass spectrometry. Analytical Biochemistry. 1999;**267**:84-91. DOI: 10.1006/abio.1998.2949
- [57] Kobayasi Y, Orga Y, Suzuki KT. Speciation and metabolism of selenium injected with <sup>82</sup>Se-enriched selenite and selenite in rats. Journal of Chromatography B: Biomedical Sciences and Applications. 2001;**760**:73-81. DOI: 10.1016/S0378-4347(01)00252-3
- [58] Duncan EG, Maher WA, Jagtap R, Krikowa F, Roper MM, O'Sullivan CA. Selenium speciation in wheat grain varies in the presence of nitrogen and sulphur fertilizers. Environmental Geochemistry and Health. 2017;**39**:955-966. DOI: 10.1007/s10653-016-9857-6
- [59] Lavu RVS, Laing GD, Van De Wiele T, Pratti VL, Willekens K, Vandecasteele B, et al. Fertilizing soil with selenium fertilizers: Impact on concentration, speciation, and bioaccessibility of selenium in leek (*Allium ampeloprasum*). Journal of Agricultural and Food Chemistry. 2012;**60**: 10930-10935. DOI: 10.1021/jf302931z
- [60] Garousi F, Domokos-Szabolesy F, Jánószky M, Kovács AB, Veres S, Soós A, et al. Selenoamino acid-enriched green pea as a value-added plant protein source for humans and livestock. Plant Foods for Human Nutrition. 2017;**73**:168-175. DOI: 10.1007/s11130-017-0606-5
- [61] Ni Z, Liu Y, Qu M. Determination of 5 natural selenium species in selenium-enriched bamboo shoots using LC-ICP-MS. Food Science and Biotechnology. 2014;**23**:1049-1053. DOI: 10.1007/s10068-014-0143-z
- [62] Gomes da Silva E, Mataveli LRV, Arruda MAZ. Speciation analysis of selenium in plankton, Brazil nut and human urine samples by HPLC-ICP-MS. Talanta. 2013;**110**:53-57. DOI: 10.1016/j.talanta.2013.02.014
- [63] Michalska-Kacymirow M, Kurek E, Smolis A, Wierzbicka M, Bulska E. Biological and chemical investigation of *Allium cepa* L. response to selenium inorganic compounds. Analytical and Bioanalytical Chemistry. 2014;**406**:3717-3722. DOI: 10.1007/s00216-014-7742-7

- [64] Ruszczyńska A, Konopka A, Kurek E, Elguera JCT, Bulska E. Investigation of biotransformation of selenium in plants using spectrometric methods. *Spectrochimica Acta Part B*. 2017;**130**:7-16. DOI: 10.1016/j.sab.2017.02.004
- [65] Larsen EH, Sloth J, Hansen M, Moesgaard S. Selenium speciation and isotope composition in <sup>77</sup>Se enriched yeast using gradient elution HPLC separation and ICP-dynamic reaction cell-MS. *Journal of Analytical Atomic Spectrometry*. 2003;**18**:310-316. DOI: 10.1039/B209586F
- [66] Khanam Am Platel K. Bioaccessibility of selenium, selenomethionine and selenocysteine from foods and influence of heat processing on the same. *Food Chemistry*. 2016;**184**:1293-1299. DOI: 10.1016/j.foodchem.2015.09.005
- [67] Zhang HJ, Gao PF, Guo XF, Wang H. Simultaneous determination of selenium containing amino acids and their sulfur-analogues in green tea and gynostemma pentaphyllum infusion with high performance liquid chromatography based on fluorescence labeling. *Microchemical Journal*. 2013;**110**:192-197. DOI: 10.1016/j.microc.2013.03.016
- [68] Tie M, Li B, Zhuang X, Han J, Liu L, Hu Y, et al. Selenium speciation in soybean by high performance liquid chromatography coupled to electrospray ionization-tandem mass spectrometry (HPLC-ESI-MS/MS). *Microchemical Journal*. 2015;**123**:70-75. DOI: 10.1016/j.microc.2015.05.017
- [69] Maneetong S, Chookhampaeng S, Chantiratikul A, Chinrasri O, Thosaikham W, Sittipout R, et al. Hydroponic cultivation of selenium-enriched kale (*Brassica oleracea* var. *alboglabra* L.) seedling and speciation of selenium with HPLC-ICP-MS. *Microchemical Journal*. 2013;**108**:87-91. DOI: 10.1016/j.microc.2013.01.003
- [70] Vonderheide AP, Wrobel K, Kannamkumarath SS, B'Hymer C, Montes-Bayon M, de Leon CP, et al. Characterization of selenium species in Brazil nuts by HPLC-ICP-MS and ES-MS. *Journal of Agricultural and Food Chemistry*. 2002;**50**:5722-5728. DOI: 10.1021/jf0256541
- [71] Marchante-Gayon JM, Thomas C, Feldmann I, Jakubowski N. Comparison of different nebulizers and chromatographic techniques for the separation of selenium in nutritional commercial supplements by hexapole collision and reaction cell ICP-MS. *Journal of Analytical Atomic Spectrometry*. 2000;**15**:1093-1102. DOI: 10.1039/B002372H
- [72] Zheng J, Kosmus W. Retention study of inorganic and organic selenium compounds on a silica-based reversed phase column with ion-pairing reagents. *Chromatographia*. 2000;**51**:338-344. DOI: 10.1007/BF02490613
- [73] Garcia-Alvarez-Coque MC, Ruiz-Angel MJ, Berthod A, Carda-Broch S. On the use of ionic liquids as mobile phase additives in high-performance liquid chromatography. Review *Analytica Chimica Acta*. 2015;**883**:1-21. DOI: 10.1016/j.aca.2015.03.042
- [74] Grijalba AC, Fiorentini EF, Wuilloud RG. Ionic liquid-assisted separation and determination of selenium species in food and beverage samples by liquid chromatography coupled to hydride generation atomic fluorescence spectrometry. *Journal of Chromatography. A*. 2017;**1491**:227-125. DOI: 10.1016/j.chroma.2017.02.045
- [75] McCalley DV. Understanding and manipulating the separation in hydrophilic interaction liquid chromatography—A review. *Journal of*



Chromatography. A. 2017;**1523**:49-71.  
DOI: 10.1016/j.chroma.2017.06.026

[76] Far J, Preud'homme H, Lobinski R.  
Detection and identification of  
hydrophilic selenium compounds in  
selenium-rich yeast by size exclusion-  
microbore normal-phase HPLC with  
the on-line ICP-MS and electrospray  
Q-TOF-MS detection. *Analytica  
Chimica Acta*. 2010;**657**:175-190. DOI:  
10.1016/j.aca.2009.10.040

[77] Dernovics M, Far J, Lobinski R.  
Identification of anionic selenium  
species in Se-rich yeast by electrospray  
QTOF MS/MS and hybrid linear ion  
trap/orbitrap MSn. *Metallomics*.  
2009;**1**:317-329. DOI: 10.1039/b901184f

[78] Greco G, Letzel T. Main interactions  
and influences of the chromatographic  
parameters in HILIC separation.  
*Journal of Chromatographic Science*.  
2013;**51**:684-693. DOI: 10.1093/  
chromsci/bmt015

[79] Sentkowska A, Pyrzynska K.  
Hydrophilic interaction liquid  
chromatography in the speciation  
analysis of selenium. *Journal of  
Chromatography B*. 2018;**1074-1075**:8-  
15. DOI: 10.1016/j.jchromb.2018.01.006

[80] Arnaudguilhem C, Bierla K,  
Ouerdane L, Preud'homme H,  
Yiannikouris Y, Lobinski R. Selenium  
metabolomics in yeast using  
complementary reversed-phase/  
hydrophilic ion interaction (HILIC)  
liquid chromatography-electrospray  
hybrid quadrupole trap/orbitrap mass  
spectrometry. *Analytica Chimica  
Acta*. 2012;**757**:26-38. DOI: 10.1016/j.  
aca.2012.10.029

[81] Bierła K, Godin S, Lobiński R,  
Szpunar J. Advances in electrospray  
mass spectrometry for the selenium  
speciation: Focus on Se-rich yeast.  
*TRAC Trends in Analytical Chemistry*.  
2018;**104**:87-94